

WHAT TO EXPECT POST YOUR URETEROSCOPY PROCEDURE

This information explains what you may experience post your procedure and how to take care of yourself during this time.

Expect:

- Sensation of needing to urinate (settles 24hrs in 50% of people)
- Some mild burning or blood in the urine (may come and go until after the stent is out)
- Stent insertion you are likely to need a stent (a fine plastic tube). It allows urine to drain from your kidney into your bladder when you have a blockage (such as a stone or growth) in your ureter.
- A further procedure to remove your stent (internal drain)
- Return to work when you are feeling comfortable. Dr Elmes rooms can provide you with a medical certificate if required.

What to do:

- · Drink plenty of fluid
- Do NOT dehydrate
- · You can exercise and do most normal activities with a stent in


Avoid:

Working/strenuous activity for 24hours

Inform Dr Elmes' rooms if:

- Unable to urinate
- Heavy continuous bleeding / multiple clots
- Fevers / unwell
- Severe stent pain (especially if with every urination)
- Bladder spasms (may need <u>Oxytrol patches</u> to settle this)

Stent insertion:



50% of people do not realise they are present; the other 50% may experience frequency, urgency or perineal/ penile tip referred pain. Use oxytrol patches for these symptoms.

A small portion (10%) will suffer from severe stent pain. **Use endone as required or mechanically obstruct your stent.** To do this, push hand in towards ribs on stent side where the pain is present and lean over on to this side to mechanically kink the stent.